REPORT DOCUMENTATION PAGE

Form Approved OMB No. 0704-0188

The public reporting burden for this collection of information is astimated to average 1 hour par response, including the time for reviewing instructions, searching existing data sources,

gathering and maintaining the data needed, and complating and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for raducing the burden, to the Department of Defense, Executive Services and Communications Directorate (0704-0188). Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to any penalty for failing to comply with a collection of information if it does not display a currently valid OM8									
PLEASE DO NO	T RETURN YOU	R FORM TO TH	E ABOVE ORGANIZATION	ON.					
	29-07-2010 2. REPORT TYPE Conference Proceeding			3. DATES COVERED (From - To)					
4. TITLE AND	SUBTITLE	-			5a. COI	NTRACT NUMBER			
Using Coupled	Models to Str	ady the Effects	of River Discharge on						
Biogeochemical Cycling and Hypoxia in the Northern Gulf of Mexico						5b. GRANT NUMBER			
					5c. PROGRAM ELEMENT NUMBER				
					N/A				
6. AUTHOR(S)					5d. PROJECT NUMBER				
			hrter, J. Hagy, B. Seha	effer, M.	NNH07CA881				
Murrell, J. Ku	rtz, B. Hercher	iroder, R. Gree	n and P. Eldridge		5e. TASK NUMBER				
					5f. WORK UNIT NUMBER				
ľ					73-9364-09-5				
7. PERFORMIN	G ORGANIZATI	ON NAME(S) A	ND ADDRESS(ES)		B. PERFORMING ORGANIZATION				
Naval Researe	h Laboratory					REPORT NUMBER			
Oceanography						NRL/PP/733009-9348			
Stennis Space	Center, MS 39	529-5004							
						40. CDOMCODIMONITORIC ACRONIVATION			
9. SPONSORIN	IG/MONITORING	AGENCY NAM	IE(S) AND ADDRESS(ES)			10. SPONSOR/MONITOR'S ACRONYM(S)			
NASA Headqu	uartare					NASA			
Attn: Laurie F						11. SPONSOR/MONITOR'S REPORT			
Mail Code 210		Rm. N111				NUMBER(S)			
Greenbelt, MD									
12. DISTRIBUTI	ON/AVAILABILI	TY STATEMENT	Г						
Approved for public release, distribution is unlimited.									
13. SUPPLEMENTARY NOTES									
14 ADSTRACT									
14. ABSTRACT We describe em		es to understand	physical processes and b	iogeochemical	cycles in o	coastal waters through the use of satellites,			
						ty to model ecological systems and the impact of			
						cal processes responsible for hypoxic events ocesses leading to hypoxia. Our efforts			
characterizes the	e carbon cycle a	ssociated with riv	er plumes and the export	of organic matt	er and nu	trients form coastal Louisiana wetlands and			
embayments in	a spatially and te	mporally intensive	e manner previously not possible to the second seco	oossible. River	ine nutrier	nts clearly affect ecosystems in the northern Gulf			
of Mexico as evidenced in the occurrence of regional hypoxia events. Less known and largely unquantified is the export of organic matter and nutrients from the large areas of disappearing coastal wetlands and large embayments adjacent to the Louisiana Continental Shelf. This project									
provides new methods to track the river plume along the shelf and to estimate the rate of export of suspended inorganic and organic particulate matter									
and dissolved organic matter form coastal habitats of south Louisiana.									
15. SUBJECT TERMS									
Culf of Maying higher and a pureles counted model									
Gulf of Mexico, biogeochemical cycles, coupled model									
16. SECURITY	CLASSIFICATIO	N OF:	17. LIMITATION OF	1B. NUMBER	19a. NA	ME OF RESPONSIBLE PERSON			
		c. THIS PAGE	ABSTRACT	OF	Bradley Penta				
Unclassified	Unclassified	Unclassified	UL	PAGES 19b. TELEPHONE NUMBER (Include area code)					
Officiassified Officiassified			7		(228) 688-4736				

PUBLICATION OR PRESEN	Pubkey: 6299 NRLINST 5600.2									
, REFERENCES AND ENCLOSURES	2. TYPE OF PUBLICATION OR PRESENTATI		3. ADMINISTRATIVE INFORMATION							
Ref: (a) NRL Instruction 5600.2 (b) NRL Instruction 5510.40D Encl: (1) Two copies of subject paper (or abstract)	() Book () Book cl () Conference Proceedings (X) Conference () Invited speaker () Multime () Journal article (refereed) () Journal	t only, not published napter ence Proceedings fereed) dia report article (not refereed) esentation, not published	STRN NRLPP/7330-09-9348 Route Sheet No. 7330/ Job Order No. 73-9364-09-5 Classification X U C Sponsor NASA approval obtained yes X no							
4. AUTHOR										
Title of Paper or Presentation										
•	ffects of River Discharge on Biogechemical Cy	cling and Hypoxia in t	he Northern Gulf of Mexico							
Author(s) Name(s) (First,MI,Last), Co	de, Affiliation if not NRL									
Bradley Penta, Richard W. Gould, B. Herchenroder, R. Green, P. M.	Robert A Arnone, R. Greene, J. Lehrter, Ja	imes Hagy, B. Schae	effer, M. Murrell, J. Kurtz,							
B. Helchemodel, R. Gleen, F. M.	. Liuriuge									
It is intended to offer this paper to the Oceans '09 MTS/IEEE Conference										
	(IVa	me of Conference)								
26-OCT - 29-OCT-09, Biloxi, MS		Conference)								
(Date, Place and Classification of Conference)										
and/or for publication in Oceans '09 MTS/IEEE Conference, Unclassified (Name and Classification of Publication) (Name of Publisher)										
After presentation or publication, pertinent publication/presentation data will be entered in the publications data base, in accordance with reference (a). It is the opinion of the author that the subject paper (is) (is notX) classified, in accordance with reference (b). This paper does not violate any disclosure of trade secrets or suggestions of outside individuals or concerns which have been communicated to the Laboratory in confidence. This paper (does) (does notX) contain any militarily critical technology. This subject paper (has) (has neverX) been incorporated in an official NRL Report.										
	dley Penta, 7331 Code (Principal Author)	mary	(Signature)							
5. ROUTING/APPROVAL	Experience of the second secon									
CODE	SIGNATURE	DATE	COMMENTS							
Author(s) Penta	Bradlyfond		Need by 11 5-p 09 Publicly accessible sources used for this publication							
	^		09-1226-2266							
Section Head Shulman	ash		09-1226-2266 abstract approval							
Branch Head Robert A Arnone, 7330	Kalin									
Division Head Ruth H. Preller, 7300	Par H Ban	8128127	Release of this paper is approved. To the best knowledge of this Division, the subject matter of this paper (has) (has neverX_) been classified.							
Security, Code			Paper or abstract was released. A security filed in this office.							
1226 Office of Counsel,Code	ABeede,		2. A copy is filed in this office.							
1008.3 ADOR/Director NCST	varieque,	9/2/2009	invention memu							

Division, Code Author, Code

E. R. Franchi; 7000

Public Affairs (Unclassified/ Unlimited Only), Code 7030.4

9/1109

Shannonbuland

PUBLICATION OR PRESENTATION RELEASE REQUEST 9. 132 6 8 NRLINST 38002								
REFERENCES AND ENCLOSURES	2. TYPE OF PUBLICATION OR PRESENT	ATION	3, ADMINISTRATIVE INFORMATION					
Ref: (a) NRL Instruction 5600,2 (b) NRL Instruction 5510,40D Enct; (1) Two Opies of subject paper (or abstract)	() Book () Book () Book () Conference Proceedings (X) Cor (refereed) (nx () Invited speaker () Multi () Journal writide (refereed) () Jou	tract only, not published ik-chapter iference Proceedings of refereed) timedia report mai article (not refereed) I Presentation, not published	STRN NRLPP/7330-69-9348 Route Sheet No. 7330/ Job Order No. 73-9364-09-5 Classification X U C Sponsor NASA approval obtained yes X no					
AUTHOR	6							
Title of Paper or Presentation								
	ffects of River Discharge on Biogechemica	l Cycling and Hypoxia in th	e Northern Gulf of Mexico					
Author(s) Nizime(s) (First,MI,Last), Code, Affilietion if not NRL Bradley Points, Richard W. Gould, Robert A Amone, R. Greene, J. Lehrter, James Hagy, B. Schaeffer, M. Murrell, J. Kurtz, B. Hercherhroder, R. Green, P. M. Eldridge								
It is intended to offer this paper to th	Oceans '09 MTS/EEE Conference	*						
26- OCT - 29- OCT- 09, Blioxi, MS		(Name of Conference)						
	(Date, Place and Classification	of Conference)						
and/or for publication in Oceans '09 MTS/IEEE Conference, Unclassified (Name and Classification of Publication) (Name of Publisher) After presentation or publication, pertinent publication/presentation data will be entered in the publications data base, in accordance with reference (a). It is the opinion of the author that the subject paper (is) (is notX) classified, in accordance with reference (b). This paper does not violate any disclosure of trade secrets or suggestions of outside individuals or concerns which have been communicated to the Laboratory in confidence. This paper (does) (does notX) contain any militarity critical technology. This subject paper (has) (has never _X) been incorporated in an official NRL Report.								
	dley Penta, 7331	Bradly	Bylas					
	Code (Principal Author)		(Signeture)					
ROUTING/APPROVAL	SIGNATURE	DATE	COMMENTS					
CODE	SIGNATURE	DATE	Need by 11 5409					
Author(*) Penta	Bralloff		Need by 11 340					
			Publicly accessible sources used for this publication					
			09-1226- 2266					
	00 0		09-1226-2266 abstract approval					
Section Head Shulman	Bh		Spoode Approval Atichel					
Branch Head Robert A Arnone, 7330	Klich							
Division Head			Release of this paper is approved. To the best knowledge of this Division, the					
Ruth H. Preller, 7300	Day H Bur	8121107	subject matter of this paper (hes) (has neverX_) been classified.					
Security, Code 1226	month benala	03/109	Paper or abstract was released. A copy is filed in this office,					
Office of Counsel,Code 1008,3	Becke	9/2/2009	invention memo					
ADOR/Director NCST E. R. Franchi; 7000	77005		This is a Final Security Review. Any changes made in the document					
Public Affairs (Unclassified/ Unlimited Only), Code 7030.4	ShannonBreland	9/1/09	nullify the Security Review					
Division, Code								
Author, Code			0110.01 4110.150					

Using Coupled Models to Study the Effects of River Discharge on Biogeochemical Cycling and Hypoxia in the Northern Gulf Of Mexico

B. Penta¹, D. Ko¹ R. Gould¹, R. Arnone¹, R. Greene², J. Lehrter², J. Hagy², B. Schaeffer², M. Murrell², J. Kurtz², B. Herchenroder³, R. Green¹, and P. Eldridge⁴

'Naval Research Laboratory Stennis Space Center, MS 39529

²USEPA/NHEERL Gulf Ecology Division 1 Sabine Island Drive Gulf Breeze, FL 32561

³Lockheed Martin Information Technology EPA Environmental Modeling and Visualization Lab. Research Triangle Park, NC 27711

⁴USEPA Coastal Ecology Branch, Western Ecology Division 2111 SE Marine Science Center Drive, Newport, OR 97365

Abstract- We describe emerging capabilities to understand physical processes and biogcochemical cycles in coastal waters through the use of satellites, numerical models, and ship observations. Emerging capabilities provide significantly improved ability to model ecological systems and the impact of environmental management actions on them. The complex interaction of physical and biogcochemical processes responsible for hypoxic events requires an integrated approach to research, monitoring, and modeling in order to fully define the processes leading to hypoxia. Our effort characterizes the carbon cycle associated with river plumes and the export of organic matter and nutrients from coastal Louisiana wetlands and embayments in a spatially and temporally intensive manner previously not possible. Riverine nutrients clearly affect ecosystems in the northern Gulf of Mexico as evidenced in the occurrence of regional hypoxia events. Less known and largely unquantified is the export of organic matter and nutrients from the large areas of disappearing coastal wetlands and large embayments adjacent to the Louisiana Continental Sheif. This project provides new methods to track the river plume along the shelf and to estimate the rate of export of suspended inorganic and organic particulate matter and dissolved organic matter from coastal habitats of south Louisiana.

I. INTRODUCTION

Along the Louisiana coast, the annual spring to summer episodes of bottom water hypoxia ($[O_2] < 2 \text{ mg } \Gamma^1$) cover an areal extent on the order of 15,000 - 20,000 km² (Fig. 1). This region of the northern Gulf of Mexico is strongly influenced by outflow from the Mississippi and Atchafalaya rivers and the many estuaries and embayments that constitute the coastal wetlands of the Mississippi River watershed. This outflow earries dissolved inorganic nutrients, particulate and dissolved organic matter, and suspended inorganic particulate matter, which, through complex interactions with the physical regime and biogeochemical processes of the region, affect the onset and duration of hypoxic events.

The goals of our project are: 1) to assess and predict the relationships between nutrient loads and hypoxia, 2) quantify the errors and uncertainties of nutrient load reduction scenarios, and 3) to provide tools to support nutrient management decisions. The complex physical and biogeochemical processes involved require an integrated approach to research, monitoring, and modeling in order to fully define the processes responsible for hypoxic events.

20100820123




Fig. 1 Bathymetry of the Louisiana Shelf (m) with the hypoxia region depicted in gray. The four regions used in the 'box model' version of the biogeochemical model (see lext) are also shown (1-4). This is the domain of the physical and coupled biogeochemical models.

II. DATA

A. In situ Sampling

The EPA Gulf Hypoxia Monitoring and Modeling Program has collected data during offshore and inshore surveys on the Louisiana shelf and coastal embayments (Barataria, Terrebonne, and Atchafalaya Bays). In situ biological, optical, physical, and chemical measurements were collected during multiple cruises over several years. Measurements included: primary productivity; water-column and sediment respiration; sediment nutrient and dissolved gas fluxes; water-column nutrients; chlorophyll a; high performance liquid chromatography (HPLC) pigment analysis; particulate organic carbon and nitrogen (POC and PON); dissolved oxygen; dissolved inorganic carbon (DIC); CO₂; pH; temperature; salinity; surface and underwater irradiance; ADCP horizontal and vertical current velocities; and in situ optical measurements (absorption and scattering coefficients and partitioned filter pad absorptions).

B. Satellite

The Louisiana coastal region comprises various environments - river plumes, streams, marshes, swamps, wetlands, and shallow embayments. These sources of water, nutrients, and inorganic and organic matter impart a complex array of optical signatures to the shelf. Techniques employed for the analysis and interpretation of satellite-derived time-series data allow us to distinguish these signatures and thus investigate the sources and fates of different water masses [1]. Multi-spectral algorithms allow for the extension of remote sensing capabilities to coastal ocean (case 2) waters where the optical signal is not necessarily chlorophyll-dominated. Calibrations of satellite sensors with accurate atmospheric corrections and ground-truthing with *in situ* measurements were used to develop a quasi-analytical algorithm [2] to separate unique optical components (Fig. 2) of absorption (phytoplankton and non-living material (CDOM + detritus)) and spectral backscattering. These components have also been used to characterize distributions of particulate organic and inorganic matter (POM, PIM) [3], both of which are important in shallow river-dominated regions where sediment resuspension plays an important role.

Ternary plots [4] of the components of total absorption are used to 'fingerprint' water masses using surface ocean color imagery. These techniques provide the capability to define the seasonal cycles of the water masses, residence times, and dispersion rates in the surface waters. Green and Gould [5] developed a multiple linear regression model with ocean color time series (2002-2004) to identify the importance of various physical forcing mechanisms, such as discharge and wind stress, in determining optical variability on the shelf. Riverinc nitrate concentration and wind stress were found to best correlate with the phytoplankton absorption in nearshore waters, indicating that stratification and vertical mixing are of primary importance to phytoplankton dynamics in the hypoxic areas. Further offshore, solar radiation and sea surface temperature (SST) were the dominant components, suggestive of upwelled nutrient sources. These satellite-derived products are also used for development and evaluation of our model system.




Fig.2 Satellite-derived (SeaWIFs) products: Weekly composites (week of June 18, 2003) of chlorophyll (OC4), TSS (total suspended solids), CDOM (gelbstuff), and absorption of phytoplankton (A_{ph}).

III. MODELS

A. Physical Model

Our model domain encompasses the area from 88 to 95 degrees West Longitude and 27 to 30.5 degrees North Latitude (Fig. 1). Over this domain we run a high-resolution physical circulation model, with 1/64-degree horizontal resolution and 36 layers in the vertical (with 19 sigma layers on the continental shelf). The model is forced with high-resolution surface fields (wind, air pressure, and shortwave radiation) from the Coupled Ocean-Atmosphere Mesoscale Prediction System (COAMPSTM). This high-resolution domain receives boundary information from the larger scale NRL IASNFS (Naval Research Laboratory Intra-Americas Seas Nowcast/ Forecast System) [6] model with tidal forcing and assimilation of satellite altimetry and sea-surface temperature and salinity (SST/SSS). The discharge of 103 rivers and river outlets are included.

The model does an excellent job of simulating the observed temperature structure (Fig. 3). In the freshest inshore waters, the model is less adept at simulating the observed salinity. We are exploring methods for adjusting the vertical distribution of freshwater input as a function of river flow-rate in order to improve the inshore salinity in the model.




Fig. 3 Regression of modeled temperature (°C; left) and salinity (PSU; right) vs. in situ temperature and salinity over four cruise transects.

B. Biogeochemical Model

We employed the Eldridge and Roelke [7] biogeochemical 'box model' in four regions along the salinity gradient off the mouth of the Mississippi River (Fig. 1); the regions are roughly defined as salinity 0-18 PSU, 18-27 PSU, 27-32 PSU, and 32-34.5 PSU. Each region (1-4) in the model consists of a surface mixed-layer, four bottom water layers, and a sediment layer (Fig. 4). The surface layer exchanges O₂ and CO₂ with the atmosphere. Within the surface layer, multiple phytoplankton species compete for nitrogen, phosphorus, and light. Six phytoplankton types (P1-P6) are defined on a range of edibility, growth rates, and sinking rates. A cell quota model allows luxury consumption of nutrients (and nutrient starvation) by the phytoplankton. Diel zooplankton vertical migration across the pyenocline allows nighttime grazing in the mixed layer that is influenced by prey edibility and stoichiometry. The zooplanton then migrate downward, to avoid predation in the sunlit waters, and excrete and respire in the sub-pyenocline waters. The sub-pyenocline aggregate water-column metabolism is also influenced by sinking fecal pellets and dead (or dying) algal cells. A simplified version of the multi-element sediment diagenesis model of Eldridge and Morse [8] is included in the bottom sediment layer. In addition, CDOM has also been added to the model (not shown). This model formulation predicts the temporal onset and duration of hypoxic and anoxic events in the four regions, but cannot (in this form) characterize the spatial extent of hypoxia.




Fig. 4 Schematic diagram of the 'box model' version of the biogeochemical model.

C. Coupled Model

By coupling the circulation and ecological models, we provide a new, improved management tool to simulate the impact of various environmental scenarios (changing discharge, nutrient, stratification, etc...) on the extent, development, maintenance, and dissipation of hypoxic events. The 'box model' has been altered such that all of the water column (mixed layer and sub-pyenoeline) processes are included in each of the 19 sigma layers of the physical model. The same advection-diffusion equations are used along with output from the physical model (offline coupling) of the three-dimensional currents (u, v, w) and the vertical mixing coefficient (k_b) to drive the biogeochemistry. The coupled model was first used to advect and diffuse inert tracer (no

biogeochemical reactions) input from the rivers and bays (Fig. 5).




Fig. 5 The physical model (white arrows represent currents) with inert tracer input from rivers (colors).

We have recently begun running the coupled model with the water column biogeochemical reactions. To validate the functionality and internal consistency of the model, an initial test simulation was run for 2003. The results indicate that the model performs in an ecologically stable manner; e.g., phytoplankton bloom and decline within reasonable concentrations and time frames. The surface chlorophyll concentration of the model is computed from the sum of all six phytoplankton (algae) groups (cell numbers times internal nutrient concentration). A snapshot (Fig. 6) of model surface chlorophyll for June 18, 2003 can be compared to the satellite-derived (Fig. 2) chlorophyll and the *in situ* measurements (the colored circles overlain on Fig. 6) for the same day. It is evident that, while the model generates chlorophyll concentrations of the correct order of magnitude nearshore, it does not simulate the offshore gradient properly. The surface chlorophyll concentrations of the six model phytoplankton groups are shown individually in Fig. 7. Phytoplankton group 6 is responsible for most of the total surface chlorophyll signal. In the model, this group is the least edible – causing the zooplankton to shift their grazing away from this group and onto the other groups. By increasing the zooplankton grazing on group 6 (increasing edibility), we may improve the model outcome.




Fig. 6 Results for total phytoplankton chlorophyll (mg CHL m³) from our initial test run of the coupled model (x-axis is degrees longitude and y-axis is degrees North latitude). Colored circles represent in situ measurements. The date is June 18, 2003 for comparison with Fig. 2.




Fig. 7 Results for each of the phytoplankton (Algae 1-6) groups (mg Chl m³) from our initial test run of the eoupled model (June 18, 2003).

The x-axis is degrees longitude and the y-axis is degrees North latitude.

The distribution of riverine CDOM input to the shelf (Fig. 8) needs to be adjusted, but the dynamics appear sound. Suspended particulate matter [SPM (TSS)] sinks out of the model too quickly (Fig. 8) and currently there is no model mechanism for resuspension. We recognize the importance of sediment resuspension and mobility to the hypoxia problem and are implementing an empirical wave model to account for these processes.




Fig. 8 Initial test run results for CDOM and suspended particulate matter (SPM) for June 18, 2003. The x-axis is degrees longitude and the y-axis is degrees North latitude.

IV. FUTURE WORK

We continue to analyze the results from our first coupled model run. In addition to the modifications mentioned above, we are currently implementing the sediment diagenesis component of the biogeochemical model into the coupled system. Another important enhancement will be the inclusion of a more accurate light propagation scheme [9] in the model. Our irradiance measurements have shown photosynthetically active radiation (PAR) penetrating deeper into the water column (oceasionally to deep benthic layers) than the 'standard' light model replicates. A more accurate depiction of the cuphotic zone may have important effects on the hypoxia simulations.

Ultimately, the goal is for these components to become a forecasting model and management tool. In order for the tool to be useful, we need to understand and quantify the uncertainties and errors inherent in our simulations. We can then investigate various nutrient-loading scenarios and determine their impact on the area of hypoxia.

ACKNOWLEDGMENT

This project is funded by the U.S. Environmental Protection Agency.

REFERENCES

- [1] Arnone, R.A., A.M. Wood, and R.W. Gould, "The evolution of optical water mass classification," Oceanography, 17(2), pp. 14-15, 2004.
- [2] Lee, Z., K.L Carder, and R.A. Arnone., "Deriving inherent optical properties from water color: a multiband quasi-analytical algorithm for optically deep waters, "Applied Optics, 41(27), pp. 5755-5772, 2002.
- [3] Gould, R.W., R.H. Stavn, M.S. Twardowski, and G.M. Lamela, "Partitioning optical properties into organic and inorganic components from ocean color imagery," *Proceedings, Ocean Optics XVI*, Santa Fe, New Mexico, Feburary 2002.
- [4] Gould, R.W., in preparation.
- [5] Green, R.E. and R.W. Gould, "A predictive model for satellite-derived phytoplankton absorption over the Louisiana shelf hypoxic zone: effects of nutrients and physical forcing. *Journal of Geophysical Research* 113, C06005, doi:10.1029/2007JC004594, 2008.
- [6] Ko, D.S., R.H. Preller, and P.J. Martin, "An Experimental Real-Time Intra Americas Sea Ocean Nowcast/Forecast System for Coastal Prediction", Proceedings, AMS 5th Conference on Coastal Atmospheric & Oceanic Prediction & Processes, pp. 97-100, 2003.
- [7] Eldridge, P.M, and D.L. Roelke, "Origins and scales of hypoxia on the Louisiana shelf: importance of seasonal plankton dynamics and river nutrients and discharge," *Ecological Modeling*, in press.
- [8] Eldridge, P.M., J.W. Morse, "Origins and temporal scales of hypoxia on the Louisiana shelf: importance of benthic and sub-pycnocline water metabolism," Marine Chemistry 108, pp. 159-171, 2008.
- [9] Penta, B., Z. Lee, R. M. Kudcla, S. L. Palacios, D. J. Gray, J. K. Jolliff, and I. G. Shulman, "An underwater light attenuation scheme for marine ecosystem models," Optics Express 21, pp. 16,581-16,591, 2008.